

HOMOMORPHISM, PARTIAL HOMOMORPHISM AND PAIR HOMOMORPHISM IN BE-ALGEBRAS

GAUTAM KUMAR PANDEY, MD. ILYAS

Department of Mathematics, Gaya College, Gaya

AND

R.L. PRASAD

Department of Mathematics, Magadh University, Bodh Gaya

RECEIVED : 30 December, 2017

Here we have discussed some homomorphisms in BE-algebras. Concept of partial homomorphism and pair homomorphism in BE-algebras have been developed with suitable examples and properties.

KEYWORDS : BE-algebra, homomorphism, partial homomorphism, pair homomorphism.

INTRODUCTION

HS. Kim and Y. N. Kim [1] introduced the concept of BE – algebra in 2006. Since then several related concepts have been developed and studied.

Definition (1.1): A BE-algebra is a system $(X; *, 1)$ consisting of a non-empty set X , a binary operation “ $*$ ” and a fixed element 1 satisfying the following conditions :-

1. (BE 1) $x * x = 1$
2. (BE 2) $x * 1 = 1$
3. (BE 3) $1 * x = x$
4. (BE 4) $x * (y * z) = y * (x * z)$

for all $x, y, z \in X$.

Definition (1.2): Let $(X; *, 1)$ and $(Y; o, e)$ be BE-algebras and let $f: X \rightarrow Y$ be a mapping. Then f is called a homomorphism if $f(x * y) = f(x) o f(y)$ for all $x, y \in X$.

SOME HOMOMORPHISMS

The following result have been established in [3, 2017].

Theorem (2.1): Let S be an universe and let X be the set of all fuzzy sets defined on S . Let 1^* and 0^* be the fuzzy sets defined on S as

$$1^*(x) = 1 \quad \text{and} \quad 0^*(x) = 0 \quad \text{for all } x \in S.$$

Also for $f, g \in X$, we define $f = g$ iff $f(x) = g(x)$ for all $x \in S$.

For $f, g \in X$ a binary operation “ o ” is defined as

$$(f o g)(x) = \min \{f(x), g(x)\} + 1 - f(x)$$

$$= (f \wedge g)(x) + 1 - f(x).$$

Then $(X; o, 1^*)$ is a BE-algebra with zero element 0^* .

Note (2.2) : For $\alpha \in [0, 1]$, the constant fuzzy set $\alpha(t) = \alpha$ for all $t \in S$ is identified as α .

Definition (2.3) : Let x be a fixed element of S . For $\delta \in X$. Let $\delta(x) = \alpha$. We consider the constant mapping $\alpha : S \rightarrow X$ defined as $\alpha(t) = \alpha$ for all $t \in S$. Let $f_x : X \rightarrow X$ be defined as

$$f_x(\delta) = \alpha = \delta(x). \tag{2.1}$$

We prove that

Theorem (2.4): f_x is a homomorphism on X for every $x \in S$.

Proof : Let $\delta_1, \delta_2 \in X$ and let $\delta_1(x) = \alpha, \delta_2(x) = \beta$. Then

$$f_x(\delta_1) = \alpha, f_x(\delta_2) = \beta.$$

Let $f_x(\delta_1 \circ \delta_2) = \gamma$. Then

$$\begin{aligned} \gamma(x) &= (\delta_1 \circ \delta_2)(x) \\ &= \min \{ \delta_1(x), \delta_2(x) \} + 1 - \delta_1(x) \\ &= \min \{ \alpha, \beta \} + 1 - \alpha \\ &= 1 \text{ or } \beta + 1 - \alpha \end{aligned}$$

according as $\alpha \leq \beta$ or $\beta < \alpha$.

Table 2.1

o	S	A	B	C	D	E	F	0
S	S	A	B	C	D	E	F	0
A	S	S	S	D	D	D	S	D
B	S	F	S	D	D	E	F	E
C	S	F	S	S	S	F	F	F
D	S	A	B	B	S	F	F	A
E	S	B	B	B	S	S	S	B
F	S	B	B	C	D	D	S	C
0	S	S	S	S	S	S	S	S

This means that

$$f_x(\delta_1 \circ \delta_2) = 1^* \text{ or } \beta + 1 - \alpha. \tag{2.2}$$

according as $\alpha \leq \beta$ or $\beta < \alpha$.

Again, $f_x(\delta_1) \circ f_x(\delta_2) = \alpha \circ \beta = 1^* \text{ or } \beta + 1 - \alpha. \tag{2.3}$

according as $\alpha \leq \beta$ or $\beta < \alpha$.

From (2.2) and (2.3) it follows that

$$f_x(\delta_1 \circ \delta_2) = f_x(\delta_1) \circ f_x(\delta_2),$$

Hence f_x is a homomorphism.

Example (2.5) : Let $S = \{a, b, c, d, e\}$ and $X = \{\phi, A, B, C, D, E, F, S\}$

where $A = \{a, b\}$, $B = \{a, b, c\}$, $C = \{c\}$, $D = \{c, d, e\}$, $E = \{d, e\}$, $F = \{a, b, d, e\}$; $S \equiv 1$, $\phi \equiv 0$.

For $L, M \in X$, we define a binary operation ‘ o ’ as

$$L o M = L^c \cup M.$$

Then Cayley table for this operation ‘ o ’ be given by

Then $(X; o, S)$ is a BE–algebra with zero element 0 [5].

For $a \in S$, let $g_a : X \rightarrow X$ be a mapping defined as $g_a(L) =$ smallest element of X containing L and a .

$$\begin{aligned} \text{Then, } g_a(A) &= A, g_a(B) = B, g_a(C) = B, g_a(D) = S, \\ g_a(E) &= F, g_a(F) = F, g_a(S) = S, g_a(0) = A. \end{aligned}$$

Now we see that

$$\begin{aligned} g_a(A o B) &= g_a(S) = S, g_a(A) o g_a(B) = A o B = S; \\ g_a(A o D) &= g_a(D) = S, g_a(A) o g_a(D) = A o S = S; \\ g_a(C o E) &= g_a(F) = F, g_a(C) o g_a(E) = B o F = F; \\ g_a(D o 0) &= g_a(A) = A, g_a(D) o g_a(0) = S o A = A; \\ g_a(E o 0) &= g_a(B) = B, g_a(E) o g_a(0) = F o A = B. \end{aligned}$$

For other elements it can be proved that

$$g_a(L o M) = g_a(L) o g_a(M) \text{ for } L, M \in X.$$

So g_a is a homomorphism.

PARTIAL HOMOMORPHISM

Definition (3.1) : Let $(X; *, 1)$ be a BE–algebra and let $f : X \rightarrow X$. If there exists a subalgebra M of X such that

$$f(x * y) = f(x) * f(y)$$

for all $x, y \in M$ but $f(x * y) \neq f(x) * f(y)$ for some $x, y \in X$, then we say that f is a partial homomorphism on X with respect to subalgebra M .

Example (3.2) : Let (X, T) be a topological space where T contains only clopen (closed and open) subsets. For $A, B \in T$ we define a binary operation ‘ o ’ on T as

$$A o B = A^c \cup B. \quad \dots (3.1)$$

Then $(T; o, 1)$ is a BE–algebra with zero element $0 \equiv \phi$ and $1 \equiv X$ [5].

For a fixed $a \in X$, let S be the collection of those elements of T which contain a . Now $L, M \in S \Rightarrow L o M = L^c \cup M \in S$, since $L^c \cup M$ contains a . So S is a BE–subalgebra of T .

Let $f_a : T \rightarrow T$ be a mapping defined as

$$f_a(L) = L \text{ or } L^c \text{ according } a \in L \text{ or } a \notin L.$$

Let $L, M \in S$, we have

$$f_a(L o M) = f_a(L^c \cup M) = L^c \cup M,$$

since $a \in M$. Also in this case

$$f_a(L) = L, f_a(M) = M \text{ and } f_a(L) \circ f_a(M) = L \circ M = L^c \cup M.$$

Thus $f_a(L \circ M) = f_a(L) \circ f_a(M)$ is satisfied for all $L, M \in S$.

Again, $a \notin L$ and $a \in M \Rightarrow f_a(L \circ M) = f_a(L^c \cup M) = L^c \cup M$, since $a \in L^c \cup M$.

In this case, $f_a(L) = L^c$ and $f_a(M) = M$ which gives $f_a(L) \circ f_a(M)$
 $= L^c \circ M = L \cup M.$

So $f_a(L \circ M) \neq f_a(L) \circ f_a(M)$ for some $L, M \in T$.

This means that f_a is a partial homomorphism in T with respect to subalgebra S .

PAIR HOMOMORPHISM

Definition (4.1) : Let $(X; *, 1)$ and $(Y; \circ, e)$ be BE-algebras.

Let $f : X \rightarrow Y$ and $g : X \rightarrow Y$ be mappings. The pair (f, g) is said to be pair homomorphism if

$$f(x * y) = g(x) \circ g(y) \quad \dots (4.1)$$

for all $x, y \in X$.

Example (4.2): In the above definition, let $f(x) = e$ for all $x \in X$ and $g(t) = a \in Y$ for all $t \in X$, then the pair (f, g) is pair homomorphism. For,

$$f(x * y) = e = g(x) \circ g(y)$$

for all $x, y \in X$.

Note (4.3) : Pair (f, g) is pair homomorphism

\Rightarrow Pair (g, f) is pair homomorphism.

In the above example $g(x * y) = a \neq e = f(x) \circ f(y)$.

Note (4.4) : If $g(1) = e$ then f coincides with g and f is a homomorphism. For $x \in X$ we have

$$\begin{aligned} f(x) &= f(1 * x) = g(1) \circ g(x) \\ &= e \circ g(x) \\ &= g(x). \end{aligned}$$

Note (4.5) : From the above example it also follows that in pair (f, g) g may not be unique for a mapping f .

Example (4.6) : Let $(X; *, 1)$ be a BE-algebra with zero element 0 and let $Y = X \times X$.

Then $(Y; \odot, (1, 1))$ is a BE-algebra [4, 2014] with zero element $(0, 0)$ where \odot is defined as

$$(x_1, x_2) \odot (y_1, y_2) = (x_1 * y_1, x_2 * y_2).$$

We consider mappings $P_1, Q_1 : Y \rightarrow Y$

defined as $P_1(x_1, x_2) = (1, x_2)$ and $Q_1(x_1, x_2) = (0, x_2)$ for all $(x_1, x_2) \in Y$. Then

$$\begin{aligned} P_1((x_1, x_2) \odot (y_1, y_2)) &= P_1(x_1 * y_1, x_2 * y_2) \\ &= (1, x_2 * y_2) \\ &= (0, x_2) \odot (0, y_2) \end{aligned}$$

$$= Q_1(x_1, x_2) \odot Q_1(y_1, y_2)$$

for all $(x_1, x_2), (y_1, y_2) \in Y$. So (P_1, Q_1) is a pair homomorphism.

Lemma (4.7) : Let $f, g : (X; *, 1) \rightarrow (Y; o, e)$ be a mapping such that pair (f, g) is a homomorphism. Then

$$f(1) = e;$$

$$x \leq y \Rightarrow g(x) \leq g(y);$$

Proof: (a) We have $f(1) = f(1 * 1) = g(1) o g(1) = e$.

(b) Also $x \leq y \Rightarrow x * y = 1$

$$\Rightarrow f(x * y) = f(1) = e$$

$$\Rightarrow g(x) o g(y) = e$$

$$\Rightarrow g(x) \leq g(y).$$

Definition (4.8) : Let (f, g) be a pair homomorphism from a BE– algebra $(X; *, 1)$ into a BE–algebra $(Y; o, e)$. Then $\text{Ker } f$ is defined as $\text{Ker } f = \{x \in X : f(x) = e\}$.

Theorem (4.9) : Let $(X; *, 1)$ and $(Y; o, e)$ be BE–algebra and let $f, g : X \rightarrow Y$. Let (f, g) be a pair homomorphism. Then

(a) $x \in \text{Ker } f$ iff $g(x) = g(1)$;

(b) $\text{Ker } f$ is a BE–sub algebra of $(X; *, 1)$.

Proof: (a) Let (f, g) be a pair homomorphism and let $x \in \text{Ker } f$.

Then $e = f(x) = f(1 * x) = g(1) o g(x)$. This implies $g(1) \leq g(x)$.

Again from lemma (4.7) (b) it follows that

$$x \leq 1 \Rightarrow g(x) \leq g(1). \text{ So } g(x) = g(1).$$

Conversely, suppose that

$$g(x) = g(1)$$

Then $f(x) = f(1 * x) = g(1) o g(x)$

$$= g(1) o g(1)$$

$$= e.$$

So $x \in \text{Ker } f$.

(b) Let $x, y \in \text{Ker } f$. Then from above $g(x) = g(y) = g(1)$. Also $f(x * y) = g(x) o g(y) = g(1) o g(1) = e$. So $x * y \in \text{Ker } f$. This proves that $\text{Ker } f$ is a BE–algebra.

REFERENCES

1. Kim, H. S. and Kim, Y. H., On BE–algebras, *Sci. Math. Japon*, **66**, No. **1**, pp. 113–116 (2007).
2. Kim, K.H., A Note on BE–algebras, *Sci. Math. Japon*, **72**, No. **2**, pp. 127–132 (2010).
3. Pandey, Gautam Kumar, Ilyas, Md. and Prasad, R.L., A class of BE–algebras, *Acta Ciencia Indica*, Vol. **XLIII M**, No. **1**, pp. 47–53 (2017).
4. Pathak, K., Sabhapandit, P. and Chetia, B.C., Cartesian product of BE/CI–algebras with Essences and Atoms, *Acta Ciencia Indica*, Vol. **XL M**, No. **3**, pp. 271–279 (2014).
5. Rani, Rashmi, BE–algebras with zero element, *Acta Ciencia Indica* (to be published).

□