

EFFECT OF AUTOMOBILE EXHAUSTS ON SOME PHYSICO-CHEMICAL CHARACTERISTICS OF ROAD SIDE SOIL AND PIPER-BETLE CROP

R. SONAM SAMARPITA, MANOJA DAS

GIET University, Gunpur

P.K. MISRA

Ex-P.G. Department of Chemistry, Ravenshaw College, Cuttack 753003 (India)

RECEIVED : 25 December, 2020

The physico chemical characteristic of road side grown piper betle (L) plant parts and soil sample has been determined. High soil contents of toxic or total heavy metals have been observed in the proximity of road side grown as compared to plant grown piper betle (L) at 100 mL. distance away from road side. Likewise, level of total Nitrogen and phosphorous also showed variations in different plant samples depending upon the extent of automobile exhaust release along road side.

INTRODUCTION

In the last 2-3 decades concern about human health hazards associated with heavy metals in soil and plant has considerably increased (Aziad *et al.*, 1984) (Zwart and Sloof, 1989). High soil contents of toxic total heavy metals have been observed along road side (Broad ford *et al.*, 1975; Rodriguez and Rodriguez, 1982; Mehta and Barbate, 1991, Sarin 1996). Heavy metal contents in certain plants have been studied (Flemming and Parle, 1977). Similarly, the uptake of Nitrogen and Phosphorus in some crops has also been studied (Tondon 1982; Bhargava, 1995).

MATERIAL AND METHODS

Collection of Samples :

In the month of June, 2015 and 2016 soil samples and piper betle (L) adult plant sample were collected from 3 sites on main NH ((5) between Cuttack-Bhubaneswar high way, viz. 20 mt., and 100 mt. distance away from the road side. Samples were collected in polythene containers and washed with 1 per cent Nitric Acid and deionized water to avoid absorption of

heavy metals on the walls of the container and also to preserve the samples for subsequent analysis of total heavy metals, Nitrogen and Phosphorous distribution in various parts. Like wise, soil samples were also collected from the same farm land.

BIOCHEMICAL TECHNIQUES

Total Nitrogen: For total nitrogen estimation, digestion will be done according to Snell and Snell (1954) and later estimation of the digest will be done colorimetrically.

The estimation of digest will be done with Nessler's reagent of Koch and Mc. Meekin's formula Oser, (1965). Amount of nitrogen will be calculated by using a calibration curve.

Total Phosphorus: For total phosphorus, digestion will be done by modified Allens method (Allen, 1940) with metol reagent (Oser, 1965). The digest of total phosphorous will be studied at 525 nm. (Allen, 1940). The amount of phosphate will then be quantitatively determined by using a calibration curve.

Total Heavy Metals: For estimation of total heavy metals, dry samples are digested in a mixture of 60% perchloric acid and concentrated nitric acid and then it is heated to dryness. The dry residue is dissolved in ammonium citrate buffer. From this solution heavy metals can be extracted with the help of purified dithizone (Sandell, 1950). The quantities will be estimated with the help of standard and calibration curve based on the methods suggested by Sandell (1950).

RESULT AND DISCUSSION

Concentration of total heavy metals in the samples, collected from different distances along road side, is shown in Table 1. Similarly, total Nitrogen and Phosphorous contents are also shown in Table 1. Amount of total heavy metals are found more in the soil and piper beetle plant parts growing in the proximity of road side as compared to those growing at 100 mt. distance in heavy metals in road side crops is probably linked to the quantity of automobile exhaust pollutants or traffic density on the road sider

Table 1. Effect of automobile exhausts on the uptake and distribution of heavy metal (per plant) in road side piper beetle (L)

Month of study	Total Heavy Metals Level in					
	Farm soil	Stem cover µmg/plant	Stem pith shoot	Leaves	Juice	Total heavy metal
Plants grown at 20 mt. distance away from road side (Experimental site I)						
June	0.556	0.1394	0.056	0.063	0.0314	0.023
Plants grown at 50 mt. distance away from road side (Experimental site II)						
June	0.385	0.0386	0.0445	0.0469	0.0305	0.266
Plants grown at 100 mt. distance away from road side (Experimental site III)						
June	0.358	0.0318	0.0428	0.0404	0.0386	0.19924

Table 2 shows that Nitrogen and Phosphorus uptake and distribution in the piper betle plant parts of road side crop. Levels of both Nitrogen and Phosphorus do not show much variations in the samples collected from different distances away from road side. However, there is some decline in both these parameters from proximity to more distances away from road side. However, there is some decline in both these parameters from proximity to more distances away from road side.

Table 2. Some biochemical characteristics of soil and *piper-betle* plant parts of 3 study site and long side, affected by automobile exhaust pollutants

Distance away from road side	Farm soil	Steam cover	Stem pita	Leaves	Juice
mg/gm dry weight					
Total Nitrogen					
20 mt. (Experimental Site I)	0.46	19.45	20.6	27.8	33.86
50 mt. (Experimental Site II)	0.48	13.95	19.5	26.8	28.88
100 mt. (Control Site)	0.55	12.48	18.13	26.36	25.86
Total Phosphate					
20 mt. (Experimental Site I)	0.13	3.98	0.686	6.85	6.86
50 mt. (Experimental Site II)	0.18	2.69	0.528	5.93	6.46
100 mt. (Control Site)	0.25	3.66	0.513	5.14	7.58

REFERENCES

- Allen, R., J.L., The estimation of phosphorous, *Biochem. J.*, **34**, 858-865 (1940).
- Azad, A.S., Arora, B.R., Bijay and Sekhon, G.S., Nature and Extent of heavy metal pollution from industrial units in Ludhiana, *Indian J Eco.*, **Vol II(I)**, 1-5 (1984).
- Bhargava, T.N., Uptake of heavy metals by some crop plant parts from solution and plluted river water, *Ph.D.Thesis*, C.C.S.Univ., Meerut, India (1995).
- Bradford,G.R., Page, A.L., Lund, L.J. and Olmstead,W., Trace element concentrations of sewage treatment plant effluent and sludges, their interactions with solid and uptake by plants, *Journa, Environ. Qual.*, **4**, 123-127 (1973).
- Deon, J.G., Bos, F.L. and Lonenetle, K.H., Removing heavy metals from waste water, *Environ Sci. Tech.*, **6**, 518-522 (1972).
- Flemming, G.A. and Parle, P.J., Heavy metals in soil, herbage and vegetables from an industrialized area, west of Dublic City, *Jr. J. Agfri Sci.*, **16(1)**, 35-48 (1977).
- Jenne, E.A., Controls of Zn, Mn, Fe, Co, Ni, Cu concentration in soils and water, the significant role of Mn and Fe oxidized, *Amm. Chem. Soc. Adr. Chem. Ser.*, **73**, 337-387 (1968).
- Kumar, R., Studies on the effects of water pollution on certain crops with particular reference to sugar mill effluent discharge, *Ph.D. Thesis*, C.C.S.University, Meerut, India (1998).
- Mehta, B.H. and Barhate, K.D., Mineral contents of cow pea cultivated in polluted water of Bombay, *Journal Ecotaxicol, Environ, Monit.*, **1(3)**: 225-229 (1991).
- Oser, B.I., *Hawk's Physiological Chemistry*, Tata Mc. Graw Hill Pub. Co. Ltd., New Delhi (1965).

11. Rodriguez, F.M. and E.Redriguez, Lead, Cadmium levels in soil and plants near highways and their correlation with traffic density. *Environ. Pollut. Ser. Bio. Chem. Phys.*, **4(4)**, 281-290 (1982).
12. Sandel, E.B., *Colorimetric defarmination of metals*, **2nd Ed.**, Inter Science Publ., in New York, (1950).
13. Sarin, R., Chemical speciation and bioavailability of heavy metals in aquatic environment. *Ab. Proc. 83rd Ind. Sci. Congress*, Patiala (Chemistry Section Part IV), page 199 (1996).
14. Sell, F.D. and Snell, C.T., *Calorimetric Methods of Analysis*. Dvon Nostrand Co., No.4 (1954).
15. Tondon, P.K., Effect of Zn and Fe. Supply on the uptake of n.P. and K in rice, *Bangladesh J. Bot.*, **11(1)**, 7-13 (1982).
16. Zwart, D. de and Sloof, W., Toxicity of mixture of heavy metals and petrochemicals on *Xenopus Laevis*. *Bull. Environ Contam Toxicolol*, **38**, 345-351 (1987).

□